

21-7-6



1411-0228

NURSERY POLYON

Average Media Temperatures	60°F	70°F	80°F	90°F
Longevity in Months	5-6	3-4	2-3	1-2

Application Rates

Incorporation

Uniformly mix (incorporate) Custom Nursery Fertilizer into potting media at following rates using trials and grower experience to obtain optimum performance. The addition of dolomitic lime-stone, additional micronutrients or other media amendments may be needed.

Topdressing - Rates Per Container

Uniformly apply product away from the base of plant.

(gallons)	Low	Medium	High
Trade G (g)	4	4	4
1 (g)	4	6	6
2 (g)	10	14	16
3 (g)	20	24	30
5 (g)	26	32	40
7 (g)	40	48	58
10 (g)	50	60	72
15 (g)	54	66	82
20 (g)	68	84	102

GUARANTEED ANALYSIS

* Total Nitrogen (N)	21.0000%
2.4200% Ammoniacal Nitrogen	
18.5800% Urea Nitrogen	
Available Phosphate (P2O5)	7.0000%
Soluble Potash (K2O)	6.0000%
Magnesium (Mg)	0.5000%
0.250% Water Soluble Magnesium (Mg)	
Sulfur (S)	3.8000%
3.800% Combined Sulfur (S)	
Copper (Cu)	0.0500%
0.050% Water Soluble Copper (Cu)	
Iron (Fe)	6.0000%
1.670% Water Soluble Iron (Fe)	
1.360% Chelated Iron (Fe)	
Manganese (Mn)	0.0500%
0.050% Water Soluble Manganese (Mn)	
Molybdenum (Mo)	0.0010%
Zinc (Zn)	0.0500%
0.050% Water Soluble Zinc (Zn)	

Derived From: Ammonium Phosphate, Polymer Coated Urea, Sulfate of Ammonia, Sulfate of Potash, Copper Sulfate, Ferrous Sulfate, Iron EDTA, Iron Humate, Iron Oxide, Iron Sucrate, Magnesium Oxide, Magnesium Sulfate, Manganese Sulfate, Sodium Molybdate, Zinc Sulfate

* 19.307% slow release NITROGEN derived from Polymer Coated Urea

Warning: --- This fertilizer is to be used only on soils which respond to Molybdenum. Crops high in Molybdenum are toxic to ruminants.

Chlorine (Cl), Not more than..... .1200%

Density - 62 lb./cu. ft.)

Directions for Use

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

CONVERSION TABLE

Rounded Measure	Gram:
1 teaspoon (tsp)	7.1
1 tablespoon (tbs.)	22.0
1/4 cup	61.4
1/2 cup	141.5

†The gram weights are approximate. Actual weight in your measuring should be determined for the fertilizer being applied.

Harrell's SPOON SIZE/RATE

Spoon Size	Grams
1	10
2	15
3	20
4	26
5	38
6	54
7	70
8	88

KEEP OUT OF REACH OF CHILDREN
CAUTION
HARMFUL IF SWALLOWED
AVOID INHALATION OF PRODUCT

APPLICATION PRECAUTIONS

- A trial of this product on a limited scale application methods, and desired rates is recommended. Avoid application to plants under stress.
- Do not incorporate into media prior to steam sterilization.
- For mixes containing soil, reduce application rate by approximately 25%.
- When using supplemental liquid feed, reduce the rate of this formulation accordingly.
- If mixed media is not used within two weeks, leach thoroughly before using.
- To avoid buildup of soluble salts, occasional leaching may be necessary.

Net Wt.
50 LBS.
(22.7 KG)

MANUFACTURED FOR HARRELL'S LLC. (F352) 720 Kraft Road, Lakeland, FL 33815 - (863) 687-2774 - (800) 282-8007
Distributed by: Harrell's, LLC. 151 Gene E. Stewart Ct., Sylacauga, AL 35151 - (256) 245-5559 - (800) 282-8007

The GREEN color of the Polyon® capsule and RLC (Reactive Layers Coating) are trademarks of RLC Technologies, L.L.C. POLYON® and POLYON NPK® are registered trademarks of RLC Technologies.

1411-0228 061411